

The Monetary Value of Disability-Adjusted Life Years in Kenya in 2017

SAGE Open
 October-December 2020: 1–15
 © The Author(s) 2020
 DOI: 10.1177/2158244020970556
 journals.sagepub.com/home/sgo


Rose Nabi Deborah Karimi Muthuri¹
 and Joses Muthuri Kirigia² 

Abstract

All diseases and injuries exert a heavy epidemiological and socioeconomic burden on Kenya. The specific objectives of this study were (a) to estimate the monetary value of disability-adjusted life years (DALYs) from all causes in 2017 with the ongoing implementation of United Nations sustainable development goal (SDG) 3 and (b) to estimate the reductions in the monetary value of DALYs in Kenya, assuming diseases and injuries related to SDG3 targets are achieved by 2030. A variant of human capital (or net output) analytical framework was applied to value DALYs from 162 diseases and injuries into their monetary equivalents. The 17.9 million DALYs from 162 causes in Kenya in 2017 had a total monetary value of Int\$ 67,012,790,388. Approximately 57% of the monetary value of DALYs emanated from communicable, maternal, neonatal, and nutritional diseases; 36% from noncommunicable diseases; and 7% from injuries. We estimate that attainment of the five SDG3 targets would lead to an Int\$ 21.5 billion (45%) decrease in the SDG3-related monetary value of DALYs by 2030. There is an urgent need for health policy-makers to use this kind of evidence when advocating among public and private sectors for increased spending on health development.

Keywords

communicable diseases, disability-adjusted life years, gross domestic product, Kenya, noncommunicable diseases, value of life

Introduction

Kenya is one of the six partner states of the East African Community (EAC) regional intergovernmental organization (EAC, 2019). It has a population of 49.4 million people, that is, 27.6% of the EAC population (International Monetary Fund [IMF], 2019). The country had a total gross domestic product (GDP) of International Dollars (Int\$) 192.3 billion and a GDP per capita of Int\$3,896.

In 2017, Kenya had a total of 17,856,955 disability-adjusted life years (DALY) from all causes (Global Burden of Disease Collaborative Network, 2018). Out of which, 57% resulted from communicable, maternal, neonatal, and nutritional diseases (CMNND); 36% from noncommunicable diseases (NCDs); and 7% from injuries. Figure 1 portrays the DALYs from all causes by 23 age groups.

The substantial loss of DALYs in Kenya could be due to two systemic weaknesses. First, the inefficiently performing national health system. Kenya's health system consists of a total of 6,655 health facilities, which includes 3,384 (50.8%) owned by the government and 3,271 (49.2%) by the private sector (World Health Organization [WHO], 2017b). The country has a health workforce density of 2.0 medical doctors per 10,000 population, and 15.4 nursing and midwifery personnel per 10,000 population. Kenya's health workforce

densities are lower than the global health workforce density of 15.1 medical doctors per 10,000 population; and 34.8 nursing and midwifery personnel per 10,000 population (WHO, 2019b).

Masters et al. (2014) facility survey of pharmaceutical availability across levels of care in Kenya found 18% of essential medicines were out of stock in referral hospitals, 26% in district hospitals, 29% in subdistrict hospitals, 33% in health centers, and 39% in dispensaries.

In 2016, per capita total current health expenditure (CHE) on health in Kenya was US\$ 66 (WHO, 2019a). Approximately US\$ 24 per capita was from domestic general government health expenditure; US\$ 29.4 per capita from domestic private health expenditure; and US\$ 12.9 per capita from external health expenditure. Kenya CHE was below the range of US\$ 76 (minimum) and US\$ 342 (maximum) per person per year of health systems investment

¹University of Pretoria, South Africa

²African Sustainable Development Research Consortium, Nairobi, Kenya

Corresponding Author:

Joses Muthuri Kirigia, Professor, African Sustainable Development Research Consortium, P.O. Box 6994 00100 GPO, Nairobi, Kenya.
 Email: muthurijoses68@gmail.com



Creative Commons CC BY: This article is distributed under the terms of the Creative Commons Attribution 4.0 License (<https://creativecommons.org/licenses/by/4.0/>) which permits any use, reproduction and distribution of

the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).




Figure 1. The disability-adjusted life years (DALYs) from all causes by age groups in Kenya in 2017.

Note. The Early neonatal = 0–6 days; late neonatal = 7–27 days; and postneonatal = 28–364 days (Institute for Health Metrics and Evaluation, 2018).

recommended for attaining health sustainable development goal (SDG) 3 (Stenberg et al., 2017).

Second, there is a challenge of suboptimal population coverage of systems that provide services for addressing social determinants of health, for example, education, food and nutrition, housing, water, sanitation, and physical security. Due to inadequate nutrition, the prevalence of stunting and wasting in children under 5 years old was 26.2% and 4.2%, respectively (WHO, 2019b). About 21.3% of adults (ages 15 and older) are illiterate (United Nations Development Programme [UNDP], 2018). About 5.3% of households live in slums (Republic of Kenya, 2013). About 41.5% of the population uses water unprotected from outside contamination, and 70% do not use improved sanitation facilities (UNDP, 2018). In the perceptions of individual well-being survey, 54% of women and 64% of men indicated they felt safe; the homicide rate is 4.9 per 100,000 people (UNDP, 2018).

As a result of under-investment in the national health system, Kenya's universal health coverage (UHC) index was 57%, implying essential health services (reproductive, maternal, newborn and child health; infectious diseases; NCDs) coverage gap of 43% (WHO, 2019b). This gap calls for unremitting evidence-based advocacy within the Kenyan government, the private sector, and the external development partners to increase investments into the national health system and the other systems that address social determinants of health.

Some economically developed (Lee et al., 2019; Schofield et al., 2019; Scholten et al., 2014; Tachkov et al., 2017) and

developing countries (Kirigia et al., 2017; Kirigia & Mwabu, 2018a) used monetary value of DALY evidence to make a case for increased investment into the health system and health programs interventions. To date, no study in Kenya has attempted to estimate the total monetary value of DALYs from all causes and among all the age groups.

The specific objectives of this study were as follows:

- to estimate the monetary value of DALYs from all causes in 2017 with the ongoing implementation of the United Nations [UN] SDG3 on health (UN, 2015);
- to appraise the monetary value of DALYs from SDG3-related diseases and injuries targets in 2030;
- to calculate the reductions in the monetary value of DALYs in Kenya, assuming diseases and injuries related to SDG3 targets are achieved by 2030.

Methods

Study Site and Disease Categories

The analysis focussed on DALY estimates for all (162) diseases in Kenya in 2017 (Global Burden of Disease Collaborative Network, 2018). These diseases included 39 CMNND; 106 NCDs, which consisted of 30 neoplasms (cancers), 11 cardiovascular diseases, 5 (five) chronic respiratory diseases, 10 digestive diseases, 7 (seven) neurological disorders, 10 mental disorders, 2 (two) substance abuse, 3 (three) diabetes and kidney diseases, 12 skin and




Figure 2. The disability-adjusted life years (DALYs) in Kenya by disease categories in 2017.

subcutaneous diseases, 3 (three) sense organ diseases, 6 (six) musculoskeletal disorders, 7 (seven) other NCDs; and 17 injuries. Figure 2 shows the DALYs in Kenya by disease categories.

Murray (1994) defines the DALY as “. . . an indicator of the time lived with a disability and the time lost due to premature mortality” (p. 441). We used DALY estimates for Kenya from the Global Burden of Disease (GBD) 2017 study (Global Burden of Disease Collaborative Network, 2018).

As the ministry of finance policy-makers and the majority of private sector Chief Executive Officers are not public health specialists, they have a challenge understanding public health indices such as the DALY. The dearth of understanding of such an index puts the health sector policy-makers at a disadvantage when advocating for sustaining (or increasing) resource allocations to health as opposed to the so-called “economically productive” sectors (World Health Organization Regional Office for Africa [WHO/AFRO], 2006). Therefore, this study attempts to convert the DALYs into their monetary equivalents to enable health sector policy-makers to use the language that those who control public and private sector budgets understand.

SDG3 Disease-Related Targets

Table 1 contains the disease-related SDG3 targets and expected percentage reductions for Kenya, assuming these

targets are attained by the end of 2030. These percentages were used in estimating the monetary value of DALYs losses that would be averted following the implementation of interventions that would lead to the achievement of SDG3 targets 3.1, 3.2, 3.3, 3.4, and 3.6.

Human Capital Approach/Net Output Analytical Framework

The Organisation for Economic Co-operation and Development (OECD, 2001, p. 18) defines human capital as “The knowledge, skills, competencies, and attributes (including physical, emotional and mental health) embodied in individuals that facilitate the creation of personal, social and economic well-being.” Premature mortality and nonfatal disability from any disease or injury erodes the health stock of Kenyans, and hence, their human capital.

More than 300 years ago, Sir William Petty pioneered the application of the human capital approach (HCA) in valuing human lives (Petty, 1699). Since then, Fein (1958), Mushkin and Collings (1959), Weisbrod (1971), Landefeld and Seskin (1982), Chisholm et al. (2010), and WHO (2009) further refined the theoretical underpinning of the HCA.

We adapted the variant of human capital (or net output) analytical framework developed and applied in the monetary valuation of DALYs among 15- to 59-year olds (Kirigia & Mwabu, 2018a) and the elderly (Kirigia et al., 2017) in

Table 1. Health SDG3 Targets and Expected Percentage Reductions for Kenya.

Target	Description	Percentage reductions in SDG targets
SDG3.1	By 2030, reduce the global maternal mortality ratio to less than 70 per 100,000 live births (United Nations [UN], 2015; WHO, 2019b). Average Annual Rate of Reduction (AARR) formula (UNICEF, 2007).	86.27% (AARR = 13.225%)
SDG3.2	By 2030, end preventable deaths of newborns, with all countries aiming to reduce neonatal mortality to at least as low as 12 per 1,000 live births (UN, 2015; WHO, 2019b).	42.857% (AARR = 3.9184%)
SDG3.3	By 2030, end the epidemics of AIDS, tuberculosis, malaria, and neglected tropical diseases and reduce hepatitis, water-borne diseases and other communicable diseases (UN, 2015). (a). Reduce global HIV-related deaths from 1,062,352 in 2015 to below 500,000 by 2020 (WHO, 2016a). (b). Malaria mortality rates will be reduced globally by at least 90% from 2015 to 2030 (WHO, 2015b, 2017g). (c). The number of tuberculosis (TB) deaths will be reduced by 90% from 2015 to 2030 (WHO, 2015a, 2017c). (d). Mortality due to vector-borne diseases will be reduced globally by at least 75% from 2016 to 2030 (Global Burden of Disease Collaborative Network, 2018; WHO, 2017d). (e). Globally reduce viral hepatitis B and C deaths from 1.4 million deaths reduced to less than 500,000 by 2030 (WHO, 2016b).	54.545% (AARR = 5.4762%) 90% (AARR = 15.166%) 90% (AARR = 15.166%) 75% (AARR = 9.428%)
SDG3.4	By 2030, reduce premature mortality due to NCDs by one third through prevention and treatment and promote mental health and well-being (Global Burden of Disease Collaborative Network, 2018; UN, 2015).	33.3% (AARR = 2.855%)
SDG3.6	By 2020, halve the number of global deaths and injuries from road traffic accidents (Global Burden of Disease Collaborative Network, 2018; UN, 2015; WHO, 2019b).	50% (AARR = 7.0905%)

Note. The AARR is the average relative percent decrease per year in prevalence or rate (UNICEF, 2007). SDG = sustainable development goal.

Kenya in 2015. Subsequently, the framework was employed to value the DALYs in 2015 within the Arab Maghreb Union (Muthuri et al., 2018), the Central African Economic and Monetary Community (Kirigia et al., 2019), and the EAC (Kirigia & Mwabu, 2018b). A similar framework was applied in estimating the aggregated productivity cost of DALYs in Africa in 2015 (WHO/AFRO, 2019c).

We choose to use the HCA to monetarily value DALYs due to the availability of data on per capita GDP, CHE per capita, and DALYs on all the countries in the African continent (Kirigia & Mwabu, 2018a; WHO/AFRO, 2019c).

Estimation of the Monetary Value of DALYs From All Causes in 2017

Kenya's total monetary value of DALYs ($TMVDALY_{2017}$) from 162 diseases and injuries in 2017 was obtained through a summation of the monetary value of DALY from each i th disease or injury ($MVDALY_i$). This was expressed as follows:

$$TMVDALY_{2017} = \sum_{i=1}^{n=162} (NHGPC_{Int\$} \times DALY_n) \quad (1)$$

where $\sum_{i=1}^{n=162}$ is the summation of monetary values of DALYs from the first to the one-hundred sixty-second disease (or injury); $NHGPC_{Int\$}$ is the nonhealth GDP calculated by subtracting CHE per person from the GDP per capita in Kenya;

$MVDALY_n$ is the monetary value of DALYs from disease (or injury) n ($n = 1, \dots, 162$) in Kenya; and $DALY_n$ is the number of DALYs from n th disease in Kenya (Kirigia & Mwabu, 2018a).

Estimation of the Monetary Value of DALYs From SDG3-Related Health Conditions Diseases in 2030

Kenya's monetary values of DALYs in 2030 were estimated assuming the SDG targets for maternal mortality ratio (MMR; Target 3.1); neonatal mortality (Target 3.2); HIV/AIDS, tuberculosis, malaria, neglected tropical diseases (NTDs) and viral hepatitis deaths (target 3.3); NCD mortality (target 3.4); and injury mortality (target 3.6) would be realized. We used Equation 2 to estimate the expected reduction in monetary values of DALYs from each j th SDG3-related health condition (HC):

$$MVDALY_{HCj2030} = MVDALY_{HCj2017} - \left[MVDALY_{HCj2017} \times \left(\frac{HC_{j2017} - HC_{jTARGET}}{HC_{j2017}} \right) \right] \quad (2)$$

where $MVDALY_{HCj2030}$ is the monetary value of DALYs in Kenya from j th health condition in 2030; $MVDALY_{HCj2017}$ is the monetary value of DALYs in Kenya from j th health

condition in 2017; and $HC_{jTARGET}$ is the j th health condition DALYs in 2030 if health condition target is achieved in Kenya. We have assumed that the implied percentage reductions in DALYs would be similar to mortality reductions because the SDG3 disease-related targets are couched in terms of mortality reductions (Kirigia & Mwabu, 2018b).

We illustrate below how the monetary value of DALYs from maternal disorders (MDs) in 2030 ($MVDALY_{MD2030}$) was estimated. From Equation 1, the monetary value of DALYs from MDs in 2017 ($MVDALY_{MD2017}$) was equal to Int\$ 919,956,631. The MMR for Kenya was 510 per 100,000 live births in 2017 (MDD2017), and SDG target 3.1 is to reduce the MMR to 70 per 100,000 live births by 2030 ($MDD_{T3.1}$). Therefore,

$$\begin{aligned} MVDALY_{MD2030} &= MVDALY_{MD2017} \\ &- \left[\begin{aligned} &MVDALY_{MD2017} \\ &\times \left(\frac{MDD_{2017} - MDD_{T3.1}}{MDD_{2017}} \right) \end{aligned} \right] \\ &= 919,956,631 \\ &- \left[919,956,631 \times \left(\frac{510 - 70}{510} \right) \right] \\ &= \text{Int\$}126,268,557\dots \end{aligned} \quad (3)$$

Estimation of Monetary Value of DALY Losses Averted by 2030

There are two ways for estimating potential reductions (savings) in the monetary value of DALYs following the full attainment of each of the five SDG3 targets. First, the saving in the monetary value of DALYs from j th health condition ($MVDALY_{HCj2030}$) can be obtained as the difference between the monetary value of DALYs in 2017 ($MVDALY_{HCj2017}$) and the monetary value of expected DALYs in 2030 ($MVDALY_{HCj2030}$) from the same condition (Kirigia & Mwabu, 2018b), that is,

$$MVDALY_{HCj-SAVINGS} = \left(\begin{aligned} &MVDALY_{HCj2017} \\ &- MVDALY_{HCj2030} \end{aligned} \right) \dots \quad (4)$$

For example, the potential reduction/savings in the monetary value of DALYs from MDs ($MVDALY_{MD-SAVINGS}$) was estimated as follows:

$MVDALY_{MD-SAVINGS} = (MVDALY_{MD2017} - MVDALY_{MD2030}) = (919,956,631 - 126,268,557) = \text{Int\$}793,688,074$. Second, the savings in the monetary value of DALYs from j th health condition ($MVDALY_{HCj2030}$) can also be obtained by multiplying $MVDALY_{HCj2017}$ by the relevant percentage reductions in SDG3 targets ($\%RHCj$) contained in Table 1. In the above example, since the $MVDALY_{MD2017} = \text{Int\$}919,956,631$ and the percentage reduction for MDs is 86.2745098%, $MVDALY_{MD-SAVINGS}$ can be calculated as follows:

$$\begin{aligned} MVDALY_{MD-SAVINGS} &= \left[\begin{aligned} &MVDALY_{MD2017} \\ &\times \left(\frac{\%RHCj}{100} \right) \end{aligned} \right] \\ &= \left[\begin{aligned} &919,956,631 \\ &\times \left(\frac{86.2745098}{100} \right) \end{aligned} \right] \\ &= \text{Int\$}793,688,074 \end{aligned} \quad (5)$$

Data Sources and Software

The DALYs data estimates for the 162 causes for Kenya were from the GBD 2017 study (Global Burden of Disease Collaborative Network, 2018); data on GDP per capita (Int\$3,896) from the IMF world economic outlook database (IMF, 2019); and data on the CHE of per person (Int\$144) from the WHO Global Health Expenditure Database (WHO, 2019a). The $NHGPC_{IntS} = \text{Int\$}3,896 - \text{Int\$}144 = \text{Int\$}3,752$. The analysis of the five equations developed in the preceding subsection was conducted using Excel Software belonging to Microsoft (New York).

Results

Estimates of the Monetary Value of DALYs in Kenya in 2017

All diseases. Kenya had a total of 17,856,955 DALYs among people of all ages in 2017 (Global Burden of Disease Collaborative Network, 2018). Those DALYs had a total monetary value of Int\$ 67,012,790,388; and an average value of Int\$ 3,753 per DALY. Approximately, 56.6% emanated from communicable, maternal, neonatal, and nutritional diseases; 35.94% from NCDs; and 7.42% from injuries.

Figure 3 displays the monetary value of DALYs from all causes by age groups. The people aged 14 years and below bore DALYs valued at Int\$ 25,964,877,620 (38.7%); the 15 to 59 years olds bore DALYs valued at Int\$ 32,888,431,641 (49.1%); and those aged above 59 years incurred DALYs valued Int\$ 8,159,481,127 (12.2%).

Communicable, maternal, neonatal, and nutritional diseases. Figure 4 depicts the monetary value of DALYs related to some CMNND categories. Out of the total monetary value of DALYs from CMNND of Int\$37,952,637,789, 28.6% resulted from HIV/AIDS and sexually transmitted infections; 19.8% from maternal and neonatal infections; 19.0% from respiratory infections and tuberculosis; 15.5% from enteric infections; 8.2% from other infectious diseases; 4.9% from nutritional disorders; and 4.1% from NTDs and malaria. Supplemental Additional File 1 presents the monetary values of DALYs associated with individual CMNND health conditions. Five health conditions (HIV/AIDS, neonatal disorders, diarrhoeal diseases, lower respiratory infections, and




Figure 3. The monetary value of disability-adjusted life years (DALYs) from all causes by age groups in Kenya (Int\$ 2019).

tuberculosis) alone accounted for 76% of the total monetary value of DALYs from CMNND.

NCDs. Figure 5 shows the monetary values of DALYs from key categories of NCDs. Of the total of DALYs from NCDs valued at Int\$ 24,084,582,837, 14.8% were attributed to cardiovascular diseases; 11.0% to digestive diseases; 10.8% to neoplasms (cancers); 10.2% to mental disorders; 8.1% to neurological disorders; 7.3% to diabetes and kidney diseases; 6.9% to musculoskeletal disorders; 5.0% to chronic respiratory diseases; 4.7% to skin and subcutaneous diseases; 4.2% to sense organ diseases; 2.2% to substance use disorders; and 14.8% to other NCDs. Supplemental Additional File 2 contains a breakdown of the monetary value of DALYs from individual NCDs. Congenital birth defects, stroke, ischaemic heart disease, cirrhosis and other chronic liver diseases, and diabetes mellitus account for 31% of the monetary value of DALYs from NCDs.

Injuries. All forms of injuries resulted in DALYs valued at Int\$ 4,975,569,762 (see Figure 6). Out of the total monetary value of DALYs from injuries, unintentional injuries accounted for Int\$2.08 billion (42%), self-harm and interpersonal violence for Int\$1.6 billion (32%), and transport injuries for Int\$1.3 billion (26%). Seventy percent of the monetary

value of DALYs from injuries was attributed to road injuries (23%), interpersonal violence (19%), falls (12%), self-harm (10%), and exposure to mechanical forces (6%).

Monetary Value of DALYs in 2017 From Five SDG3-Related Targets

In 2017 SDG-related health conditions and diseases were associated with 12,627,303 DALY, which had a monetary value of Int\$ 47,387,208,551 (70.7%). As indicated in Table 2, NCDs, HIV/AIDS, and neonatal disorders accounted for 86.83% of the monetary value of SDG-related DALYs.

Estimates of Reductions in Monetary Value of DALY Losses in Kenya if the Five SDG3-Related Targets Are Achieved by 2030

Table 3 presents the monetary value of DALYs in 2017, the monetary value of DALYs in 2030, and envisaged monetary reductions by SDG health conditions.

We estimate that the full attainment of the five SDG3 targets would lead to a 45.42% decrease in the monetary value of DALYs by 2030.




Figure 4. The monetary value of disability-adjusted life years (DALYs) from communicable, maternal, neonatal, and nutritional diseases in Kenya (2019 Int\$).

Discussion

Key Findings

The 17,856,955 DALYs among people of all ages in 2017 had a total monetary value of Int\$ 67,012,790,388. Of these,

56.6% was attributed to CMNND, 35.94% to NCD, and 7.4% to injuries. The 15- to 59-year olds incurred close to half of the monetary value of DALYs. We estimate that the attainment of the five SDG3 targets would lead to an Int\$ 21.5 billion (45.4%) decrease in the monetary value of DALYs by 2030.




Figure 5. The monetary value of disability-adjusted life years (DALYs) from noncommunicable diseases in Kenya (Int\$ 2019).

SDG3.1 and SDG3.2: Maternal and Neonatal Disorders

Were Kenya to attain the SDG targets 3.1 and 3.2 by 2030, the country would annually prevent loss of DALYs with a monetary value of Int\$ 793,688,074. That kind of saving might be achieved if both the national road map for accelerating the attainment of the millennium development goals (MDGs) related to maternal and newborn health (Republic of Kenya, 2010c); and the reproductive, maternal, newborn, child, and adolescent health (RMNCAH) investment framework (Republic of Kenya, 2016b) is fully implemented in all the 47 Administrative Counties. The global strategy for women's, children's, and adolescents' health (2016–2030) (Every Woman Every Child, 2015); the United Nations General Assembly (UNGA) Resolutions on the girl child (UN, 2017a) and the right to food (UN, 2017c); and the World Health Assembly (WHA) Resolution on immunization (WHO, 2017f) underpinning the implementation of the national roadmap and framework to end preventable maternal and neonatal deaths.

SDG3.3: Tuberculosis, HIV/AIDS, Malaria, Acute hepatitis, and NTDs

The realization of SDG target 3.3 by 2030 would annually save Kenya DALYs with a monetary value of Int\$ 2,092,156,033 from TB, Int\$ 5,711,558,233 from HIV/AIDS, Int\$982,595,151 from malaria, Int\$ 114,448,409 from acute hepatitis, and Int\$335,362,674 from NTDs. In other words, the country stands to save DALYs worth Int\$7.144 billion.

The saving could be made through full implementation of Kenya's AIDS strategic framework 2014/2015 to 2018/2019 (Republic of Kenya, 2014a); the HIV and AIDS research agenda 2014/15 to 2018/19 (Republic of Kenya, 2014b); the national strategic plan for tuberculosis, leprosy, and lung health 2019 to 2023 (Republic of Kenya, 2018f); the HIV and AIDS/STI and TB multisectoral strategic plan and implementation framework 2015 to 2020 (EAC, 2015); the malaria strategy 2019 to 2023 (Republic of Kenya, 2018a); the malaria communication strategy 2016–21 (Ministry of Health, 2016); the guidelines for the treatment of chronic hepatitis B and C viral infections




Figure 6. The monetary value of disability-adjusted life years (DALYs) from injuries in Kenya (Int\$ 2019).

Table 2. Monetary Value of DALYs in 2017 From Five SDG3-Related Targets (2019 Int\$).

SDG3 targets	DALYs in 2017 ^a	Monetary value of DALYs (Int\$ or PPP) ^b	Percent ^c
SDG3.1: Maternal disorders	245,142	919,956,631	1.9
SDG3.2: Neonatal disorders	1,755,556	6,588,173,277	14.0
SDG3.3a: HIV mortality	2,790,267	10,471,190,093	22.1
SDG3.3b: Malaria mortality	290,926	1,091,772,390	2.3
SDG3.3c: TB deaths	619,443	2,324,617,815	4.9
SDG3.3d: NTDs	119,153	447,150,232	0.9
SDG3.3e: Acute hepatitis	47,440	178,030,859	0.4
SDG3.4: NCDs	6,417,832	24,084,582,837	50.8
SDG3.6: Traffic injuries	341,544	1,281,734,417	2.7
Total for SDG-related	12,627,303	47,387,208,551	100
National total	17,856,955	67,012,790,388	

Note. DALY = disability-adjusted life years; SDG = sustainable development goal; PPP = purchasing power parity.

^aDALYs data from IHME database (Global Burden of Disease Collaborative Network, 2018). ^bMonetary value of DALYs (Int\$ or PPP). ^cPercent are authors estimates.

(Ministry of Health, 2014a); the national strategic plan for control of NTDs 2016 to 2020 (Republic of Kenya, 2015a); and the national health promotion strategy (Ministry of Health, 2012).

A pertinent legal framework, including the Health Act (Republic of Kenya, 2017a), the Public Health Act (Republic of Kenya, 2012g), the HIV and AIDS prevention and control Act (Republic of Kenya, 2012d), the sexual offenses Act (Republic of Kenya, 2012h), and the malaria prevention Act (Republic of Kenya, 2012f) buttresses the implementation of communicable diseases strategies and plans.

At the global level, the UNGA declared political commitments for the fight against HIV/AIDS (UN, 2016b), malaria (UN, 2018a), TB (UN, 2018c), and NTDs (UN, 2017b; WHO, 2017e), which provide high-level political backing for the implementation of pertinent national policies, strategies, and plans.

SDG3.4: NCDs

If Kenya reduces the epidemiological burden from NCDs by one-third by 2030 (SDG3 target 3.4), it will save DALYs

Table 3. Monetary Value of DALYs by SDG Health Conditions.

Health condition/diseases	(A). Monetary value of DALYs in 2017 (Int\$)	(B). Monetary value of DALYs in 2030 (Int\$)	(C). Reduction in monetary value of DALYs in 2030 (Int\$) [C=A-B]
SDG3.1: Maternal disorders	919,956,631	126,268,557	793,688,074
SDG3.2: Neonatal disorders	6,588,173,277	3,764,670,444	2,823,502,833
SDG3.3: Tuberculosis	2,324,617,815	232,461,781	2,092,156,033
SDG3.3: HIV/AIDS	10,471,190,093	4,759,631,861	5,711,558,233
SDG3.3: Malaria	1,091,772,390	109,177,239	982,595,151
SDG3.3: Acute hepatitis	178,030,859	63,582,449	114,448,409
SDG3.3: NTDs (a+b..+l)	447,150,232	111,787,558	335,362,674
(a). Schistosomiasis	220,314,394	55,078,598	165,235,795
(b). Intestinal nematode infections	55,333,984	13,833,496	41,500,488
(c). Leishmaniasis	44,498,104	11,124,526	33,373,578
(d). Cysticercosis	35,821,824	8,955,456	26,866,368
(e). Rabies	25,760,440	6,440,110	19,320,330
(f). Dengue	14,136,229	3,534,057	10,602,172
(g). Yellow fever	12,394,580	3,098,645	9,295,935
(h). Trachoma	3,781,714	945,429	2,836,286
(i). Lymphatic filariasis	2,320,028	580,007	1,740,021
(j). Cystic echinococcosis	559,598	139,899	419,698
(k). Leprosy	197,041	49,260	147,781
(l). Other neglected tropical diseases	32,032,296	8,008,074	24,024,222
SDG3.4: Non-communicable diseases (a+b+c+...+l)	24,084,582,837	16,056,468,840	8,028,113,997
(a). Neoplasms	2,596,076,616	1,730,726,397	865,350,218
(b). Cardiovascular diseases	3,567,563,123	2,378,387,307	1,189,175,816
(c) Chronic respiratory diseases	1,205,075,880	803,387,937	401,687,943
(d). Digestive diseases	2,639,735,063	1,759,832,175	879,902,889
(e). Neurological disorders	1,962,721,321	1,308,487,423	654,233,898
(f). Mental disorders	2,463,518,824	1,642,354,095	821,164,730
(g). Substance use disorders	521,794,946	347,865,036	173,929,909
(h). Diabetes and kidney diseases	1,759,422,548	1,172,954,230	586,468,318
(i). Skin and subcutaneous diseases	1,139,408,497	759,609,463	379,799,034
(j). Sense organ diseases	1,020,302,380	680,204,988	340,097,392
(k). Musculoskeletal disorders	1,656,194,793	1,104,135,383	552,059,410
(l). Other noncommunicable diseases	3,552,768,846	2,368,524,406	1,184,244,439
SDG3.6: Transport injuries	1,281,734,417	640,867,208	640,867,208
Total (Int\$)	47,387,208,550	25,864,915,938	21,522,292,613

Note. DALY = disability-adjusted life years; SDG = sustainable development goal.

worth Int\$ 8,028,113,997. The saving is achievable through full implementation of the national health policies, strategic plans, and guidelines. For instance, the Kenya national health policy 2014 to 2030 (the Republic of Kenya, 2014b); the Kenya health sector strategic and investment plan (KHSSIP) (Republic of Kenya, 2014c); the national strategy for the prevention and control of NCD 2015 to 2020 (Republic of Kenya, 2015b); the national nutrition action plan 2012 to 2017 (Republic of Kenya, 2012e); the mental health policy 2015 to 2030 (Republic of Kenya, 2016a); the national cancer control strategy 2017 to 2022 (Republic of Kenya, 2017b); the cancer screening guidelines (Republic of Kenya, 2018b); the guidelines for cardiovascular diseases management (Republic of Kenya, 2018c); the national diabetes strategy 2010 to 2015 (Republic of Kenya, 2010b); and the

physical activity action plan (Republic of Kenya, 2018e) plus the guidelines for healthy diets and physical activity (Republic of Kenya, 2017c).

The WHA resolutions on physical activity (WHO, 2018c); infant and child feeding (WHO, 2018a); rheumatic fever and rheumatic heart disease (WHO, 2018b); cancer prevention and control (WHO, 2017a); global strategy for the prevention and control of NCD (WHO, 2000); and action plan for the prevention and control of NCDs (WHO, 2013) informed the development of Kenyan policies and strategic documents.

Various national legal instruments, such as the Kenya Constitution (Republic of Kenya, 2010a), the cancer prevention and control Act (Republic of Kenya, 2012c), the tobacco control Act (Republic of Kenya, 2012i), and the alcoholic

drinks control Act (Republic of Kenya, 2012a) undergird the implementation of NCD prevention and control strategies and plans.

At the global level, in 2018 and 2012, the UNGA committed to providing strategic leadership through whole-of-government and health-in-all-policies approaches for scale-up of the prevention and control of NCDs (UN, 2012, 2018d).

SDG3.6: Transport Injuries

In case Kenya succeeded in reducing by half the epidemiological burden from road accidents, it would reduce DALYs with a monetary value of Int\$640,867,208. The country developed the policy on integrated national transport (Republic of Kenya, 2009), the strategy and action plan for road safety (Republic of Kenya, 2015c), and the violence and injury prevention and control action plan 2018 to 2022 (Republic of Kenya, 2018d) to stem the tide of deaths and injuries associated with transport injuries. The national strategy on road safety consists of the following five components (Republic of Kenya, 2015c):

- Road safety management: Including road safety coordination and management, mainstreaming, funding, data systems, research, and monitoring and evaluation.
- Safer road and mobility: Encompassing development and maintenance of safer roads, and the safeguarding of vulnerable road users.
- Safer vehicles: Includes vehicle safety standards, compliance, and public transport regulations.
- Safer road users: Incorporating driver training, testing and licensing; road safety legislations and regulations; traffic law enforcement; road safety awareness/public education; and road safety, children, and education.
- Postcrash care: Involving availability and accessibility to emergency services and rehabilitation.

Kenya's national transport and safety authority, Act No. 33 of 2012, provides the legal framework for implementation of the aforementioned national policy, strategies, and plans (Republic of Kenya, 2014d).

Kenya's road safety policy framework is underpinned by the UNGA resolution A/RES/64/255 that proclaimed the period 2011 to 2020 as the Decade of Action for Road Safety (UN, 2010). The UNGA resolutions A/RES/72/271 (UN, 2018b) and A/RES/70/260 (UN, 2016a) both entitled "improving global road safety" provide further political support for sustained implementation of strategies for curbing transport-related deaths and nonfatal injuries of people.

Limitations of the Study

The study reported in this article has some limitations. First, in Kenya, civil registration of births and deaths is a national

Government function based in the Ministry of Interior and Coordination of National Government. The process hinges on a community-based system of civil registration, where assistant chiefs (local government administrators) record in register books births and deaths occurring at home. In contrast, health workers record those occurring at a health facility (Republic of Kenya, 2012b). In some cases, due to ignorance of the Government's requirement, parents might not report births and deaths that occur at home. Furthermore, even when parents (or next of kin) report deaths occurring at home, the cause of death might be wrong because often there are no inquiries as to cause of death.

For these reasons, the mortality statistics upon which years of life lost (YLL) calculations are based might not be accurate. The completeness of cause of death data in the WHO African region was about 6% between 2009 and 2017 due to weak civil registration systems in countries (WHO, 2019b). The GBD 2017 Mortality Collaborators (2018) and GBD 2017 Cause of Death Collaborators (2018) discuss limitations related to the accuracy of the mortality and causes of death data used the GBD 2017 study. The GBD 2017 Disease and Injury Incidence and Prevalence Collaborators (2018) expounds on the shortcomings of the national data used in estimating years living with a disability (YLD) for various diseases and injuries. Therefore, as explained clearly by the GBD 2017 DALYs and HALE Collaborators (2018), DALY estimates are influenced by the availability and quality of data for YLL and YLD.

Second, unlike the willingness-to-pay/willingness-to-accept approach proffered by some economists, the HCA does not take into account the strength of individual preferences to either avert or accept the risk of incurring a DALY loss (Jones-Lee, 1985; Mishan, 1971). Furthermore, if strictly applied, HCA would value DALY losses occurring among anyone below the minimum work age and past retirement age at zero (Jones-Lee, 1985).

Third, GDP as a measure of the well-being of societies has weaknesses, such as ignoring economic, social, political and ecological inequalities and their ramifications (Stiglitz et al., 2010); does not capture the quality of life, happiness, and well-being; disregards negative environmental consequences of economic production processes (Giannetti et al., 2015; Ivkovic, 2016); omitting nonmarketed services, for example, the contribution of homemakers, including child care, cooking, cleaning, washing, fetching water, hewing firewood, cultivating, inculcating societal values, caring for the elderly, and nursing the sick at home (Card & Mooney, 1977).

Finally, since the SDG3 disease-related targets are framed in terms of mortality reductions, we have assumed that the implied reductions in DALYs would be similar. Of course, the reductions in mortality may not result in equal reductions in DALYs. For example, some of the medical interventions for chronic diseases (e.g., NCDs) may improve health-related quality of life, but may not prolong the life expectancy of persons living with those conditions. Therefore, the use of

SDG3 disease-related targets may have led to an overestimation of the expected DALY reductions, and hence, expected monetary savings.

Conclusions

In 2017, the DALYs in Kenya had a monetary value equivalent to 34.8% of the national 2019 GDP. Full attainment of the disease and injuries related to SDG3 targets would lead to a 45.4% decrease in the monetary value of DALYs by 2030. There exists national health-related legislation, policies, strategies, and plans which, if fully funded and implemented, would enable Kenya to strengthen its health system and health programs, and consequently, assure UHC of essential health services related to SDG3 targets.

Therefore, there is an urgent need for health policy-makers to use the kind of evidence reported in this article to advocate among public and private sectors for increased spending on health development to the levels recommended by Stenberg et al. (2017).

Acknowledgments

R.N.D.K.M. is very grateful for the support and encouragement of her PhD thesis supervisors (Dr Flavia Senkubuge and Professor Charles Hongoro) at the School of Health Systems and Public Health, University of Pretoria, Pretoria, South Africa. The authors owe profound gratitude to Jehovah Jireh for meeting all their needs in the entire process of the study reported in this paper. The views expressed in this paper are exclusively those of the authors and not for the institutions of affiliation.

Author Contributions

R.N.D.K.M. and J.M.K. designed the study; reviewed pertinent literature; collated data from IHME, IMF and WHO databases; analyzed data, and wrote the paper. Both authors read and approved the final manuscript.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Ethical Approval and Consent to Participate

This study did not involve the use of animal or human data or products. The analysis was based solely on secondary statistical data published in the international databases. Therefore, ethical approval and consent to participate did not apply to the study reported in this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Joses Muthuri Kirigia  <https://orcid.org/0000-0002-2317-4666>

Supplemental Material

Supplemental material for this article is available online.

Availability of Data and Materials

The secondary data analyzed during this study is publicly available from the following websites:

- DALY data from IHME database (Global Burden of Disease Collaborative Network, 2018): <http://ghdx.healthdata.org/gbd-results-tool>
- GDP data from IMF database (IMF, 2019): <https://www.imf.org/external/pubs/ft/weo/2018/02/weodata/index.aspx>
- Per capita current health expenditure data from WHO Database (WHO, 2019a): <http://apps.who.int/nha/database/ViewData/Indicators/en>

References

- Card, W. I., & Mooney, G. H. (1977). What is the monetary value of human life? *British Medical Journal*, 2, 1627–1629. <https://doi.org/10.1136/bmj.2.6103.1627>
- Chisholm, D., Stanciole, A. E., Edejer, T. T. T., & Evans, D. B. (2010). Economic impact of disease and injury: Counting what matters. *British Medical Journal*, 340, 583–586. <https://www.jstor.org/stable/i40030885>
- East African Community. (2015). *HIV and AIDS/STI and TB multisectoral strategic plan and implementation framework 2015–2020*.
- East African Community. (2019, June 15). *EAC at a glance*. <https://www.eac.int/>
- Every Woman Every Child. (2015). *The global strategy for women's, children's and adolescents' health (2016-2030)*. <http://globalstrategy.everywomaneverychild.org/>
- Fein, R. (1958). *Economics of mental illness*. Basic Books.
- GBD 2017 Cause of Death Collaborators. (2018). Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980–2017: A systematic analysis for the Global Burden of Disease Study 2017. *Lancet*, 392(10159), 1736–1788. [https://doi.org/10.1016/S0140-6736\(18\)32203-7](https://doi.org/10.1016/S0140-6736(18)32203-7)
- GBD 2017 DALYs and HALE Collaborators. (2018). Global, regional, and national disability-adjusted life-years (DALYs) for 359 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990–2017: A systematic analysis for the Global Burden of Disease Study 2017. *Lancet*, 392(10159), 1859–1922. [https://doi.org/10.1016/S0140-6736\(18\)32335-3](https://doi.org/10.1016/S0140-6736(18)32335-3)
- GBD 2017 Disease and Injury Incidence and Prevalence Collaborators. (2018). Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: A systematic analysis for the Global Burden of Disease Study 2017. *Lancet*, 392(10159), 1789–1858. [https://doi.org/10.1016/S0140-6736\(18\)32279-7](https://doi.org/10.1016/S0140-6736(18)32279-7)
- GBD 2017 Mortality Collaborators. (2018). Global, regional, and national age-sex-specific mortality and life expectancy, 1950–2017: A systematic analysis for the Global Burden of Disease Study 2017. *Lancet*, 392(10159), 1684–1735. [https://doi.org/10.1016/S0140-6736\(18\)31891-9](https://doi.org/10.1016/S0140-6736(18)31891-9)

- Giannetti, B. F., Agostinho, F., Almeida, C., & Huisingsh, D. (2015). A review of limitations of GDP and alternative indices to monitor human wellbeing and to manage eco-system functionality. *Journal of Cleaner Production*, 87, 11–25.
- Global Burden of Disease Collaborative Network. (2018). *Global Burden of Disease Study 2017 (GBD 2017) results*. Institute for Health Metrics and Evaluation (IHME). <http://ghdx.healthdata.org/gbd-results-tool>
- Institute for Health Metrics and Evaluation (IHME). (2018). *GBD 2017 online tools overview*.
- International Monetary Fund. (2019, March 5). *World economic outlook database*. <https://www.imf.org/external/pubs/ft/weo/2018/02/weodata/index.aspx>
- Ivković, A. F. (2016). Limitations of the GDP as a measure of progress and well-being. *Ekonomski Vjesnik / Econviews*, XXIX, 257–272. <https://hrcak.srce.hr/ojs/index.php/ekonomski-vjesnik/article/view/4217/2459>
- Jones-Lee, M. W. (1985). The value of life and safety: A survey of recent developments. *The Geneva Papers on Risk and Insurance*, 10, 141–173. <https://link.springer.com/content/pdf/10.1057/gpp.1985.13.pdf>
- Kirigia, J. M., Mburugu, G. N., & Huka, G. S. (2017). The indirect cost of disability adjusted life years lost among the elderly in Kenya. *International Archives of Medicine*, 10(213), 1–14. <https://doi.org/10.3823/2483>
- Kirigia, J. M., Muthuri, R. N. D. K., & Gitonga, N. (2019). Fiscal value of disability-adjusted life years lost in the Economic and Monetary Community of Central Africa (CEMAC). *International Public Health Journal*, 11(2), 155–183. http://www.novapublishers.org/catalog/product_info.php?products_id=66020
- Kirigia, J. M., & Mwabu, G. M. (2018a). Haemorrhage of gross domestic product from disability-adjusted life years among 15-59 year olds in Kenya. *International Archives of Medicine*, 11(20), 1–19. <https://doi.org/10.3823/2561>
- Kirigia, J. M., & Mwabu, G. M. (2018b). The monetary value of disability-adjusted life years lost in the East African Community in 2015. *Modern Economy*, 9, 1360–1377. <https://doi.org/10.4236/me.2018.97087>
- Landefeld, J. S., & Seskin, E. P. (1982). The economic value of life: Linking theory to practice. *American Journal of Public Health*, 72(6), 555–566. <https://doi.org/10.2105/AJPH.72.6.555>
- Lee, Y.-R., Cho, B., Jo, M.-W., Ock, M., Lee, D., Kim, M. J., & Oh, I.-H. (2019). Measuring the Economic Burden of Disease and Injury in Korea, 2015. *Journal of Korean Medical Science*, 34(Suppl. 1): Article e80. <https://doi.org/10.3346/jkms.2019.34.e80>
- Masters, S. H., Burstein, R., DeCenso, B., Moore, K., Haakenstad, A., Ikilezi, G., Achan, J., Osei, I., Garshong, B., Kisia, C., Njuguna, P., Babigumira, J., Kumar, S., Hanlon, M., & Gakidou, E. (2014). Pharmaceutical availability across levels of care: Evidence from facility surveys in Ghana, Kenya, and Uganda. *PLOS ONE*, 9(12), Article e114762. <https://doi.org/10.1371/journal.pone.0114762>
- Ministry of Health. (2012). *National health promotion strategy for Kenya—Re-orienting health (2013–2018)*.
- Ministry of Health. (2014). *Guidelines for the treatment of chronic hepatitis B and C viral infections in Kenya*.
- Ministry of Health. (2016). *The Kenya malaria communication strategy 2016–2021*. National Malaria Control Programme.
- Mishan, E. J. (1971). Evaluation of life and limb: A theoretical approach. *Journal of Political Economy*, 79(4), 687–705. <https://www.journals.uchicago.edu/doi/abs/10.1086/259784>
- Murray, C. J. L. (1994). Quantifying the burden of disease: The technical basis for disability-adjusted life years. *Bulletin of the World Health Organization*, 72(3), 429–445. <https://apps.who.int/iris/handle/10665/52181>
- Mushkin, S. J., & Collings, F. A. (1959). Economic costs of disease and injury. *Public Health Reports*, 74, 795–809. <https://doi.org/10.2307/4590578>
- Muthuri, R. D. K., Muthuri, N. G., & Kirigia, J. M. (2018). Pecuniary value of disability-adjusted life years in the Arab Maghreb Union in 2015. *Journal of Human Resource and Sustainability Studies*, 6, 249–274. <https://doi.org/10.4236/jhrss.2018.64041>
- Organisation for Economic Co-operation and Development. (2001). *The well-being of nations—the role of human and social capital*.
- Petty, W. (1699). *Political arithmetic, or a discourse: Concerning the extent and value of lands, people, buildings: Husbandry, manufacture, commerce, fishery, artisans, seamen, soldiers; public revenues, interest, taxes, superlucration, registries, banks*. Robert Caluel.
- Republic of Kenya. (2009). *Integrated national transport policy: Moving a working nation*. Ministry of Transport.
- Republic of Kenya. (2010a). *The constitution of Kenya, 2010*. National Council for Law Reporting.
- Republic of Kenya. (2010b). *Kenya national diabetes strategy 2010-2015*. Ministry of Health.
- Republic of Kenya. (2010c). *National road map for accelerating the attainment of the MDGs related to maternal and newborn health in Kenya*. Ministry of Health.
- Republic of Kenya. (2012a). *Alcoholic Drinks Control Act, 2010*. National Council for Law Reporting.
- Republic of Kenya. (2012b). *Births and deaths registration ACT chapter 149*. National Council for Law Reporting.
- Republic of Kenya. (2012c). *The Cancer Prevention and Control Act, 2012*. Republic of Kenya.
- Republic of Kenya. (2012d). *HIV and AIDS Prevention and Control Act No. 14 of 2006. [Rev. 2012]*. National Council for Law Reporting.
- Republic of Kenya. (2012e). *Kenya's national nutrition action plan 2012-2017*. Ministry of Health.
- Republic of Kenya. (2012f). *Malaria prevention Act CAP. 246*. National Council for Law Reporting.
- Republic of Kenya. (2012g). *Public Health Act CAP. 242*. National Council for Law Reporting.
- Republic of Kenya. (2012h). *The Sexual Offences Act No. 3 of 2006, Act No. 7 of 2007, Act No. 6 of 2009, Act No. 8 of 2010, Act No. 12 of 2012*. National Council for Law Reporting.
- Republic of Kenya. (2012i). *Tobacco Control Act, 2007*. National Council for Law Reporting.
- Republic of Kenya. (2013). *2012/2013 Kenya national housing survey. Basic report*. Ministry of Land, Housing & Urban Development.
- Republic of Kenya. (2014a). *Kenya AIDS strategic framework 2014/2015 - 2018/2019*. Ministry of Health.

- Republic of Kenya. (2014b). *The Kenya health policy 2014-2030*. Ministry of Health.
- Republic of Kenya. (2014c). *Kenya health sector strategic and investment plan, 2013-2017*. Ministry of Health.
- Republic of Kenya. (2014d). *National Transport and Safety Authority Act No. 33 of 2012*. National Council for Law Reporting.
- Republic of Kenya. (2015a). *The 2nd Kenya national strategic plan for control of neglected tropical diseases 2016-2020*. Ministry of Health.
- Republic of Kenya. (2015b). *Kenya national strategy for the prevention and control of non-communicable diseases 2015 – 2020*. Ministry of Health.
- Republic of Kenya. (2015c). *National road safety action plan 2015 – 2020*.
- Republic of Kenya. (2016a). *Kenya mental health policy 2015 – 2030*. Ministry of Health.
- Republic of Kenya. (2016b). *Kenya reproductive, maternal, newborn, child and adolescent health (RMNCAH) investment framework*. Ministry of Health.
- Republic of Kenya. (2017a). *The Health Act No. 21 of 2017*. National Council for Law Reporting.
- Republic of Kenya. (2017b). *National cancer control strategy 2017 – 2022*. Ministry of Health.
- Republic of Kenya. (2017c). *National guidelines for healthy diets and physical activity*. Ministry of Health.
- Republic of Kenya. (2018a). *Kenya malaria strategy (KMS) 2019–2023*. Ministry of Health.
- Republic of Kenya. (2018b). *Kenya national cancer screening guidelines*. Ministry of Health.
- Republic of Kenya. (2018c). *Kenya national guidelines for cardiovascular diseases management*. Ministry of Health.
- Republic of Kenya. (2018d). *Kenya national violence and injury prevention and control action plan 2018 – 2022*. Ministry of Health.
- Republic of Kenya. (2018e). *National physical activity action plan 2018-2023*. Ministry of Health.
- Republic of Kenya. (2018f). *National strategic plan for tuberculosis, leprosy and lung health 2019-2023*. Ministry of Health.
- Schofield, D., Cunich, M., Shrestha, R., Passey, M., Veerman, L., Tanton, R., & Kelly, S. (2019). The indirect costs of ischemic heart disease through lost productive life years for Australia from 2015 to 2030: Results from a microsimulation model. *BMC Public Health*, *19*, Article 802. <https://doi.org/10.1186/s12889-019-7086-5>
- Scholten, A. C., Haagsma, J. A., Panneman, M. J. M., van Beeck, E. F., & Polinder, S. (2014). Traumatic brain injury in the Netherlands: Incidence, costs and disability adjusted life years. *PLOS ONE*, *9*(10), Article e110905. <https://doi.org/10.1371/journal.pone.0110905>
- Stenberg, K., Hanssen, O., Edejer, T. T. T., Bertram, M., Brindley, C., Meshreky, A., Rosen, J. E., Stover, J., Verboom, P., Sanders, R., & Soucat, A. (2017). Financing transformative health systems towards achievement of the health sustainable development goals: A model for projected resource needs in 67 low-income and middle-income countries. *Lancet Global Health*, *5*(9), e875–e887. [http://doi.org/10.1016/S2214-109X\(17\)30263-2](http://doi.org/10.1016/S2214-109X(17)30263-2)
- Stiglitz, J., Sen, A., & Fitoussi, J.-P. (2010). *Mis-measuring our lives: Why GDP doesn't add up*. The New Press.
- Tachkov, K., Kamusheva, M., Pencheva, V., & Mitov, K. (2017). Evaluation of the economic and social burden of chronic obstructive pulmonary disease (COPD). *Biotechnology & Biotechnological Equipment*, *31*(4), 855–861. <https://doi.org/10.1080/13102818.2017.1335616>
- UNICEF. (2007). *How to calculate average annual rate of reduction (AARR) of underweight prevalence. statistics and monitoring section/division of policy and practice technical note*.
- United Nation. (2010). *Improving global road safety* (UN General Assembly Resolution A/RES/64/255).
- United Nation. (2012). *Political declaration of the high-level meeting of the general assembly on the prevention and control of non-communicable diseases* (UN General Assembly Resolution A/RES/66/2).
- United Nation. (2015). *Transforming our world: The 2030 Agenda for Sustainable Development* (UN General Assembly Resolution A/RES/70/1).
- United Nation. (2016a). *Improving global road safety* (UN General Assembly Resolution A/RES/70/260).
- United Nation. (2016b). *Political declaration on HIV and AIDS: On the fast track to accelerating the fight against HIV and to ending the AIDS epidemic by 2030* (UN General Assembly Resolution A/RES/70/266).
- United Nation. (2017a). *The girl child* (UN General Assembly Resolution A/RES/72/154).
- United Nation. (2017b). *The human rights to safe drinking water and sanitation* (UN General Assembly Resolution A/RES/72/178).
- United Nation. (2017c). *The right to food* (UN General Assembly Resolution A/RES/72/173).
- United Nation. (2018a). *Consolidating gains and accelerating efforts to control and eliminate malaria in developing countries, particularly in Africa, by 2030* (UN General Assembly Resolution A/RES/72/309).
- United Nation. (2018b). *Improving global road safety* (UN General Assembly Resolution A/RES/72/271).
- United Nation. (2018c). *Political declaration of the high-level meeting of the General Assembly on the fight against tuberculosis* (UN General Assembly Resolution A/RES/73/3).
- United Nation. (2018d). *Political declaration of the third high-level meeting of the General Assembly on the prevention and control of non-communicable diseases* (UN General Assembly Resolution A/RES/73/2).
- United Nations Development Programme. (2018). *Human development indices and indicators: 2018 statistical update*.
- Weisbrod, B. A. (1971). Costs and benefits of medical research: A case study of poliomyelitis. *Journal of Political Economy*, *79*, 527–544. <https://doi.org/10.1086/259766>
- World Health Organization. (2000). *Global strategy for the prevention and control of noncommunicable diseases*.
- World Health Organization. (2009). *WHO guide to identifying the economic consequences of disease and injury*.
- World Health Organization. (2013). *Global action plan for the prevention and control of noncommunicable diseases 2013-2020*.
- World Health Organization. (2015a). *The end TB strategy*.
- World Health Organization. (2015b). *Global technical strategy for malaria 2016-2030*.

- World Health Organization. (2016a). *Global health sector strategy on HIV 2016-2021: Towards ending AIDS* (The Sixty-Ninth WHA Document A69/31).
- World Health Organization. (2016b). *Global health sector strategy on viral hepatitis, 2016–2021*.
- World Health Organization. (2017a). *Cancer prevention and control in the context of an integrated approach* (World Health Assembly Resolution WHA70.12).
- World Health Organization. (2017b). *Global atlas of medical devices*.
- World Health Organization. (2017c). *Global tuberculosis report 2018*.
- World Health Organization. (2017d). *Global vector control response 2017–2030*.
- World Health Organization. (2017e). *Global vector control response: An integrated approach for the control of vector-borne diseases* (World Health Assembly Resolution WHA70.16).
- World Health Organization. (2017f). *Strengthening immunization to achieve the goals of the global vaccine action plan* (World Health Assembly Resolution WHA70.14).
- World Health Organization. (2017g). *World malaria report 2017*.
- World Health Organization. (2018a). *Infant and young child feeding* (World Health Assembly Resolution WHA71.9).
- World Health Organization. (2018b). *Rheumatic fever and rheumatic heart disease* (World Health Assembly Resolution WHA71.14).
- World Health Organization. (2018c). *WHO global action plan on physical activity 2018–2030* (World Health Assembly Resolution WHA71.6).
- World Health Organization. (2019a, July 3). *Global health expenditure database*. <http://apps.who.int/nha/database/ViewData/Indicators/en>
- World Health Organization. (2019b). *World health statistics overview 2019: Monitoring health for the SDGs*. https://www.who.int/gho/publications/world_health_statistics/2019/en/
- World Health Organization Regional Office for Africa. (2006). *Health financing: A strategy for the African region* (WHO Regional Committee for Africa Resolution AFR/RC56/R5).
- World Health Organization Regional Office for Africa. (2019c). *A heavy burden: The productivity cost of illness in Africa*.